

I. Epidemiological profile

Population	2014	%
High transmission (>1 case per 1000 population)	24 200 000	100
Low transmission (0–1 cases per 1000 population)	0	0
Malaria free (0 cases)	0	0
Total	24 200 000	

Parasites and vectors

Major plasmodium species: *P. falciparum* (100%), *P. vivax* (0%)
 Major anopheles species: *An. gambiae*, *An. funestus*, *An. nili*

Programme phase: Control

Reported confirmed cases: 2 298 979 Estimated cases, 2013: [2 000 000–5 100 000]
 Reported deaths: 5714 Estimated deaths, 2013: [8900–20 000]

II. Intervention policies and strategies

Intervention	Policies/strategies	Yes/No	Adopted
ITN	ITNs/LLINs distributed free of charge	Yes	2001
	ITNs/LLINs distributed to all age groups	No	-
IRS	IRS is recommended	Yes	2003
	DDT is authorized for IRS	No	-
Larval control	Use of larval control recommended	Yes	2009
IPT	IPT used to prevent malaria during pregnancy	Yes	2005
Diagnosis	Patients of all ages should receive diagnostic test	Yes	2010
	Malaria diagnosis is free of charge in the public sector	Yes	2006
Treatment	ACT is free for all ages in public sector	Yes	2006
	Sale of oral artemisinin-based monotherapies	are allowed	-
	Single dose of primaquine is used as gametocidal medicine for <i>P. falciparum</i>	No	-
	Primaquine is used for radical treatment of <i>P. vivax</i>	Yes	2006
	G6PD test is a requirement before treatment with primaquine	Yes	2006
	Directly observed treatment with primaquine is undertaken	No	-
	System for monitoring of adverse reactions to antimalarials exists	Yes	2006
Surveillance	ACD for case investigation (reactive)	No	-
	ACD of febrile cases at community level (pro-active)	No	-
	Mass screening is undertaken	No	-
	Uncomplicated <i>P. falciparum</i> cases routinely admitted	No	-
	Uncomplicated <i>P. vivax</i> cases routinely admitted	No	-

Antimalarial treatment policy

Medicine	Adopted
First-line treatment of unconfirmed malaria	AL 2006
First-line treatment of <i>P. falciparum</i>	AL 2006
Treatment failure of <i>P. falciparum</i>	QN 2006
Treatment of severe malaria	AS; QN 2006
Treatment of <i>P. vivax</i>	-
Dosage of primaquine for radical treatment of <i>P. vivax</i>	0.25 mg/kg (14 d)
Type of RDT used	<i>P. f + P. v</i> specific (Combo)

Therapeutic efficacy tests (clinical and parasitological failure, %)


Medicine	Year	Min	Median	Max	Follow-up	No. of studies	Species
AL	2013–2013	2.7	7.2	11.7	28 days	2	<i>P. falciparum</i>
DHA-PPQ	2013–2013	0	0	0	28 days	2	<i>P. falciparum</i>

Insecticide susceptibility bioassays (reported resistance to at least one insecticide for any vector at any locality)

Year	Pyrethroid	DDT	Carbamate	Organophosphate	Species/complex tested
2010–2015	Yes	Yes	Yes	No	<i>An. coustani</i> , <i>An. gambiae s.l.</i>

III. Financing

Sources of financing




Financing by intervention in 2014


No data reported for 2014

IV. Coverage


ITN and IRS coverage




Cases tested and treated in public sector



Cases tracked




Test positivity




V. Impact

Confirmed malaria cases per 1000 and ABER



Malaria admissions and deaths



Impact: Insufficiently consistent data to assess trends